

ON SOME SUB-RIEMANNIAN OBJECTS IN HYPERSURFACES OF SUB-RIEMANNIAN MANIFOLDS

KANG-HAI TAN, XIAO-PING YANG

ABSTRACT. We study some sub-Riemannian objects (such as horizontal connectivity, horizontal connection, horizontal tangent plane, horizontal mean curvature) in hypersurfaces of sub-Riemannian manifolds. We prove that if a connected hypersurface in a contact manifold of dimension more than three is noncharacteristic or with isolated characteristic points, then given two points, there exists at least one piecewise smooth horizontal curve in this hypersurface connecting them. In any sub-Riemannian manifold, we obtain the sub-Riemannian version of the fundamental theorem of Riemannian geometry: there exists a unique nonholonomic connection which is completely determined by the sub-Riemannian structure and a complement of the horizontal bundle, is “symmetric” and is compatible with the sub-Riemannian metric. We use this nonholonomic connection to study horizontal mean curvature of hypersurfaces.

Keywords: sub-Riemannian manifolds, Contact manifolds, Chow’s condition, horizontal connectivity, horizontal connection, horizontal mean curvature

2000 Mathematics Subject Classification: 58A05, 58A30, 58A50, 58C50, 58J60.

1. INTRODUCTION

Recently there is an explosion of interest in the theory of sub-Riemannian manifolds (or Carnot-Carathéodory spaces in general), and in the ramification of this subject into analysis and geometry, see e.g. [1], [5], [6], [10], [11], [12], [13], [14], [17], [20] and [23]. We recall a sub-Riemannian manifold (M, Δ, g_c) is a smooth manifold M with a distribution Δ (a subbundle of the tangent bundle TM) which is endowed with a fiberwise inner product g_c (usually called sub-Riemannian metric). g_c is usually realized as the restriction on Δ of some Riemannian metric g on TM . Carnot groups are particularly interesting sub-Riemannian manifolds. Roughly speaking, the sub-Riemannian geometry of (M, Δ, g_c) is the geometry determined by the sub-Riemannian structure (Δ, g_c) which yields Carnot-Carathéodory distance if Δ satisfies the Chow’s condition and M is connected, and should be independent of the choice of the Riemannian metric g which is the extension of g_c to TM , although it often interacts with the Riemannian geometry of (M, g) .

It is natural to study the geometry of hypersurfaces (or submanifolds of codimension more than one) in sub-Riemannian manifolds not only from the geometric viewpoint (see [5], [14], [15] and [23] for the program of sub-Riemannian minimal surfaces in Carnot groups), but also from the viewpoint of analysis (see [10]–[13], [20] and [24] for the developing geometric measure theory in the setting of sub-Riemannian geometry). To develop a theory of sub-Riemannian minimal surfaces, a sub-Riemannian counterpart of the notion of the usual mean curvature on submanifolds should be laid down. In [23], [5] (see also [15]) an analysis definition of the notion of horizontal mean curvature for noncharacteristic hypersurfaces in Carnot groups has been given. In [27] we have formulated a geometric definition of the horizontal mean curvature by

using the notion of horizontal connection. It turns out that our definition coincides with the analysis definition in the case of Carnot groups. The geometric definition is valid in general sub-Riemannian manifolds.

In this paper we continue our study on geometry and calculus of hypersurfaces in sub-Riemannian manifolds. The notion of the horizontal tangent planes of smooth noncharacteristic (see Definition 3.1) hypersurfaces plays an important role in the story of [27]. The horizontal tangent plane $T_x^{\mathcal{H}}S$ at a point x in a smooth noncharacteristic hypersurface S is defined as the $k-1$ (assuming Δ is k dimensional) dimensional linear space such that

$$\Delta_x = n^{\mathcal{H}}(x) \bigoplus^{g_c} T_x^{\mathcal{H}}S$$

where Δ_x is the fiber of Δ through x , $n^{\mathcal{H}}(x)$ the horizontal normal of S at x and $\bigoplus^{g_c}$ denotes the orthogonal decomposition with respect to g_c . Since $n^{\mathcal{H}}(x)$ is by definition the projection onto Δ_x of the Riemannian normal $n(x)$ computed with respect to g (any orthogonal extension of g_c), we have

$$T_x^{\mathcal{H}}S = T_xS \cap \Delta_x.$$

We call

$$T^{\mathcal{H}}S := \bigcup_{x \in S} T_x^{\mathcal{H}}S$$

the horizontal tangent bundle. In general, if S possesses characteristic points, one also can define $T^{\mathcal{H}}S = TS \cap \Delta$. Note that $T^{\mathcal{H}}S$ is independent of the orthogonal extensions of g_c . A very interesting question is that under what condition $T^{\mathcal{H}}S$ satisfies the Chow's condition with respect to TS , that is, the subbundle $T^{\mathcal{H}}S$ (if it is a subbundle, say) together with all its Lie commutators spans TS (we assume Δ satisfies the Chow's condition). This question is relevant to another two: whether there exist sufficiently many horizontal curves in S and whether one can define a Carnot-Carathéodory metric with respect to $T^{\mathcal{H}}S$ with an induced metric $g_c^{\mathcal{H}}$ (from g_c)? We point out that in his seminal paper [17] Gromov discussed, among other things about geometry of hypersurfaces in sub-Riemannian manifolds, the relationship between the restricted Carnot-Carathéodory metric on S and the Carnot-Carathéodory metric induced by the structure $(T^{\mathcal{H}}S, g_c^{\mathcal{H}})$. He claimed that the two metrics are Lipschitz equivalent in some special cases, e.g. contact manifolds of dimension more than three, see page 104 and page 173 in [17]. But he did not seem to give conditions to guarantee that the Carnot-Carathéodory metric induced by the structure $(T^{\mathcal{H}}S, g_c^{\mathcal{H}})$ can be well defined. One of main results of this paper is to give the first result in this regard. We have

Theorem 1.1. *Let M be a smooth orientable contact manifold of dimension $2n+1$ with a contact form η and a Riemannian metric g and let $\Delta := \ker(\eta)$ be the canonical distribution. Let S be a smooth connected hypersurface in M . If $n > 1$ and there does not exist immersed submanifold contained in the set Σ_S of all characteristic points in S , then for any points $p, q \in S$, there exists a piecewise smooth curve $\gamma(t), t \in [a, b]$ connecting p, q such that $\dot{\gamma} \in T_{\gamma(t)}^{\mathcal{H}}S$. In particular, if S is noncharacteristic, $T^{\mathcal{H}}S$ as a subbundle of TS satisfies the Chow's condition, that is, $T^{\mathcal{H}}S$ together with all its Lie commutators span TS .*

Theorem 1.1 follows from Sussmann's famous Orbit Theorem and the contact structure of M . A trivial example shows that the condition of dimension more than three is necessary.

The next point we concentrate on is that whether there exists an intrinsic nonholonomic connection D in (M, Δ, g_c) . A natural candidate is the one $\tilde{D}$ obtained by projecting to Δ the

Levi-Civita connection ∇ with respect to some orthogonal extension g of g_c . We prove that $\tilde{D}$ is independent of the choice of orthogonal extension of g_c . Moreover we have

Theorem 1.2. *Let $\{X_1, \dots, X_k\}$ be an orthonormal basis of Δ . We define D by*

$$D_U V = \sum_{i=1}^k U(V^i) X_i + \sum_{i=1}^k \sum_{j=1}^k \sum_{l=1}^k U^j V^i \Gamma_{ij}^l X_l \text{ for any } U = \sum_{j=1}^k U^j X_j, V = \sum_{i=1}^k V^i X_i \in \Gamma(\Delta) \quad (1.1)$$

where $\Gamma_{ij}^l = -\frac{1}{2}\{g_c(X_i, [X_j, X_l]^{\mathcal{H}}) + g_c(X_j, [X_l, X_i]^{\mathcal{H}}) - g_c(X_l, [X_i, X_j]^{\mathcal{H}})\}$ and $X^{\mathcal{H}}$ is understood as the projection of X to Δ with respect to the direct summation decomposition $TM = \Delta \oplus \tilde{\Delta}$ where $\tilde{\Delta}$ is the distribution complementary to Δ . Then D is independent of the choice of orthonormal basis of Δ and it is the **unique** nonholonomic connection satisfying

- (1) $D_U V$ is R -linear in both arguments,
- (2) $D_U V$ is $C^\infty(M)$ -linear in the argument of U ,
- (3) the Leibniz rule holds:

$$D_U(fV) = (Uf)V + fD_U V \text{ for any } f \in C^\infty(M), U, V \in \Gamma(\Delta),$$

- (4) D is compatible with respect to g_c , that is,

$$Ug_c(V, W) = g_c(D_U V, W) + g_c(U, D_V W) \text{ for any } U, V, W \in \Gamma(\Delta), \quad (1.2)$$

- (5) the following “symmetry” property holds:

$$D_U V - D_V U = [U, V]^{\mathcal{H}} \text{ for any } U, V \in \Gamma(\Delta). \quad (1.3)$$

In particular we have $D = \tilde{D}$.

Theorem 1.2 is the counterpart of the existence and uniqueness of the Levi-Civita connection in Riemannian geometry. But the connection certainly depends on the splitting of TM .

We will use D to study the horizontal mean curvature of hypersurfaces in (M, Δ, g_c) . Let $D^\top$ be the tangent horizontal connection on the horizontal tangent bundle $T^{\mathcal{H}}S$ of a smooth noncharacteristic hypersurface S . We define the horizontal mean curvature of S as the trace of the horizontal fundamental second form which is by definition a bilinear map Π from $\Gamma(T^{\mathcal{H}}S) \times \Gamma(T^{\mathcal{H}}S)$ to N :

$$\Pi(X, Y) = D_X Y - D_X^\top Y$$

for any $X, Y \in \Gamma(T^{\mathcal{H}}S)$ where $\Gamma(T^{\mathcal{H}}S)$ denotes the set of all smooth sections of $T^{\mathcal{H}}S$ and N is the horizontal normal. The symmetry of Π follows from the symmetry property (1.3) of D and the definition of the horizontal tangent plane. Since both D and $T^{\mathcal{H}}S$ are intrinsic, so is the horizontal mean curvature.

The paper is organized as follows. In Section 2 we collect some facts about sub-Riemannian manifolds which will be used later, mainly to fix some notations. In Section 3 Theorem 1.1 is proven after introducing the notion of the horizontal tangent bundle. Roughly speaking, if we project Riemannian objects onto the horizontal bundle, such as Riemannian connection, normal vector and tangent bundle, then we get corresponding sub-Riemannian analogues: horizontal connection, horizontal normal vector and horizontal bundle. Theorem 1.2 is proven in Section 4.

2. BASIC MATERIAL ON SUB-RIEMANNIAN MANIFOLDS

Let M be a smooth (C^∞) manifold of dimension m endowed with a smooth distribution (called *horizontal bundle*) Δ of dimension k with $k < m$.¹ If we a priori equip Δ with an inner product g_c (called *sub-Riemannian metric*), we call (M, Δ, g_c) a *sub-Riemannian manifold* with the *sub-Riemannian structure* (Δ, g_c) . Let $\{X_1, \dots, X_k\}$ be an orthonormal local basis of Δ . A piecewise smooth curve $\gamma(t), t \in [a, b]$ in M is horizontal if $\dot{\gamma}(t) \in \Delta_{\gamma(t)}$ a.e. $t \in [a, b]$. The length $\ell(\gamma)$ of the horizontal curve $\gamma(t), t \in [a, b]$ is the integral $\int_a^b g_c(\dot{\gamma}(t), \dot{\gamma}(t)) dt$. Denote by L_i the set of all vector fields spanned by all commutators of X_j 's of order $\leq i$ and let $L_i(p)$ be the subspace of evaluations at p of all vector fields in L_i . We call Δ satisfies the *Chow's condition*² if for any $p \in M$, there exists an integer $r(p)$ such that $L_{r(p)}(p) = T_p M$ (the tangent space of M at p). If M is connected and Δ satisfies the Chow's condition, the Chow connectivity theorem asserts that there exists at least one piecewise smooth horizontal curve connecting two given points (see [4], [1] or [17]), and thus (Δ, g_c) yields a metric (called *Carnot-Carathéodory metric*) d_c by letting $d_c(p, q)$ as the infimum among the lengths of all horizontal curves joining p to q .

Δ is *equiregular* if the dimension of $L_i(p)$ does not depends on p for any i , that is, the tangent bundle TM is filtered by smooth subbundles

$$\Delta = L_1 \subset L_2 \subset \dots \subset L_r = TM$$

where r is called the degree of Δ . Note that we always can extend g_c to a Riemannian metric g in M such that TM can be g -orthogonally decomposed as $TM = \Delta \overset{g}{\oplus} \tilde{\Delta}$ where $\tilde{\Delta}$ is the distribution complementary to Δ . We call such g an *orthogonal extension* of g_c . Obviously the orthogonal extension of g_c is not unique in general. We will use $\Gamma(\Delta)$ to denote the set of all smooth sections of Δ .

Example 2.1 (Carnot groups). *The most interesting models of sub-Riemannian manifolds are Carnot groups (called also stratified groups). A Carnot group G is a connected, simply connected Lie group whose Lie algebra $\mathcal{G}$ admits the grading $\mathcal{G} = V_1 \oplus \dots \oplus V_l$, with $[V_1, V_i] = V_{i+1}$, for any $1 \leq i \leq l-1$ and $[V_1, V_l] = 0$ (the integer l is called the step of G). Let $\{e_1, \dots, e_n\}$ be a basis of $\mathcal{G}$ with $n = \sum_{i=1}^l \dim(V_i)$. Let $X_i(g) = (L_g)_* e_i$ for $i = 1, \dots, k := \dim(V_1)$ where $(L_g)_*$ is the differential of the left translation $L_g(g') = gg'$ and let $Y_i(g) = (L_g)_* e_{i+k}$ for $i = 1, \dots, n-k$. We call the system of left-invariant vector fields $\Delta := V_1 = \text{span}\{X_1, \dots, X_k\}$ the horizontal bundle of G . If we equip Δ an inner product g_c such that $\{X_1, \dots, X_k\}$ is an orthonormal basis of Δ , (G, Δ, g_c) is an equiregular sub-Riemannian manifold. In (G, Δ, g_c) , d_c is invariant with respect to left translation, that is $d_c(p_0 p, p_0 q) = d_c(p, q)$ for any $p_0, p, q \in G$, and is 1-homogeneous with respect to the natural dilations, that is $d_c(\delta_s p, \delta_s q) = s d_c(p, q)$ for any $s > 0, p, q \in G$, where $\delta_s p = \exp(\sum_{i=1}^l s^i \xi_i)$ for $p = \exp(\sum_{i=1}^l \xi_i), \xi_i \in V_i$.*

Example 2.2 (contact manifolds). *Contact manifolds are also interesting sub-Riemannian manifolds not necessarily with group structure. A contact manifold is a smooth (connected) manifold M with a contact structure which is by definition a codimension one distribution $\Delta \subset TM$ with non-degenerate curvature form $\omega : \Delta \wedge \Delta \rightarrow TM/\Delta$ which is defined as follows: first represent Δ locally as the kernel of a 1-form, say η on M , then identify TM/Δ with the trivial line bundle and finally define ω as $d\eta|_{\Delta}$. η is called a contact form of M (not unique). If M is orientable, η and ω can be globally defined. The non-degeneracy of ω makes the dimension of Δ be even and so the dimension of M is odd, say $2n + 1$. Note that the non-vanishing of ω on Δ*

¹Note that this imposes topological constraints on M , see [19]

²In the subelliptic theory, Chow's condition is also called Hörmander's condition.

makes the commutators of degree ≤ 2 span TM and thus (M, Δ, g_c) where g_c is the restriction on Δ of some Riemannian metric g on M is an equiregular sub-Riemannian manifold.

Example 2.3 (Heisenberg group). Heisenberg group $\mathbb{H}^n$ as a representative both in the class of Carnot groups and in the class of contact manifolds is of paramount importance, and is worthy of being paid more attention to. The underlying manifold of this Lie group is simply $\mathbb{R}^{2n+1}$, with the noncommutative group law

$$p \cdot p' = (x, y, t) \cdot (x', y', t') = (x + x', y + y', t + t' + 2(\langle x', y \rangle - \langle x, y' \rangle))$$

where we have let $x, x', y, y' \in \mathbb{R}^n, t, t' \in \mathbb{R}$. A simple computation shows that

$$\begin{aligned} X_j(p) &\stackrel{\text{def}}{=} (L_p)_* \left(\frac{\partial}{\partial x_j} \right) = \frac{\partial}{\partial x_j} + 2y_j \frac{\partial}{\partial t}, \quad j = 1, \dots, n, \\ X_{n+j}(p) &\stackrel{\text{def}}{=} (L_p)_* \left(\frac{\partial}{\partial y_j} \right) = \frac{\partial}{\partial y_j} - 2x_j \frac{\partial}{\partial t}, \quad j = 1, \dots, n, \\ T(p) &\stackrel{\text{def}}{=} (L_p)_* \left(\frac{\partial}{\partial t} \right) = \frac{\partial}{\partial t} \end{aligned}$$

for any $p = (x, y, t)$ in $\mathbb{H}^n$. We note that

$$[X_j, X_{n+k}] = -4T\delta_{jk}, \quad j, k = 1, \dots, n, \\ \text{and all other commutators are trivial,}$$

therefore the vector fields $X = \{X_1, \dots, X_{2n}\}$ constitute a basis of the Lie algebra $\mathfrak{h}_n = \mathbb{R}^{2n+1} = V_1 \oplus V_2$, where $V_1 = \mathbb{R}^{2n} \times \{0\}_t, V_2 = \{0\}_{x,y} \times \mathbb{R}$. Note that the horizontal bundle $\Delta = \text{span}\{X_1, \dots, X_{2n}\}$ is the kernel of the 1-form $\eta = \frac{1}{4}dt + \frac{1}{2}\sum_{i=1}^n(x_i dy_i - y_i dx_i)$ and the curvature form $\omega = d\eta = \sum_{i=1}^n dx_i \wedge dy_i$ is the standard symplectic form in $\mathbb{R}^{2n}$. Thus a smooth curve $\gamma(s) = (x(s), y(s), t(s)) : [a, b] \rightarrow \mathbb{H}^n$ is horizontal if and only if

$$2\dot{t}(s) = \sum_{i=1}^n y_i(s)\dot{x}_i(s) - x_i(s)\dot{y}_i(s) \quad \text{for any } s \in [a, b]. \quad (2.1)$$

For the theory of sub-Riemannian geodesics we refer to the book [22] and references therein, and see in particular [17] for a comprehensive treatment of the geometry (more than sub-Riemannian geodesics) in sub-Riemannian manifolds, also for many potential research directions.

3. THE HORIZONTAL TANGENT BUNDLE AND HORIZONTAL CONNECTIVITY IN HYPERSURFACES

Let (M, Δ, g_c) be a sub-Riemannian manifold. **In this section we always assume M is connected and Δ satisfies the Chow's condition.** By S we always mean a smooth hypersurface (i.e. an embedded submanifold of codimension 1) in M . Let g be any orthogonal extension of g_c .

Definition 3.1 (characteristic points). A point $p \in S$ is a *characteristic point* if $\Delta_p \subset T_p S$. Let Σ_S denote the set of all characteristic points in S . If $\Sigma_S = \emptyset$, we call S is *noncharacteristic*.

Typically S possesses characteristic points, see e.g. [6]. But we have

Proposition 3.2. Σ_S is a closed subset of S , and $\mathcal{H}^{m-1}(\Sigma_S) = 0$. Here $\mathcal{H}^{m-1}$ denotes the $m-1$ dimensional Hausdorff measure with respect to the Riemannian metric g (recall M is m -dimensional).

The closedness follows from the smoothness of S and the smallness of Σ_S is due to Derridj, see [7], [8]. In the case of Carnot groups of step two, when the smoothness of S is weaker than C^∞ , say $C^{1,1}$, the smallness of Σ_S is obtained by Magnani, see [20] or [21].

By Proposition 3.2, if $p \in S \setminus \Sigma_S$, then there exists a neighborhood $\mathcal{U}$ of p , such that $\mathcal{U} \cap S \subset S \setminus \Sigma_S$.

Definition 3.3 (horizontal normal). Let n^g denote the Riemannian normal of S with respect to g . We define the *horizontal normal* $n^{\mathcal{H}}$ of S as the projection of the Riemannian normal onto the horizontal bundle, that is,

$$n^{\mathcal{H}} = \sum_{i=1}^k g(n^g, X_i) X_i$$

where $\{X_1, \dots, X_k\}$ is an orthonormal basis of Δ .

It is easily seen that $p \in S$ is a characteristic point if and only if $n^{\mathcal{H}}(p) = 0$.

Definition 3.4 (horizontal tangent plane). For $p \in S$, we define

$$T_p^{\mathcal{H}}S := \{v \in \Delta_p : g_c(v, n^{\mathcal{H}}(p)) = 0\}$$

as the *horizontal tangent plane* of S at p .

From the definition we see that if $p \in \Sigma_S$ then $T_p^{\mathcal{H}}S = \Delta_p$, and otherwise $T_p^{\mathcal{H}}S$ is a $k-1$ dimensional subspace of Δ_p . Thus

$$T^{\mathcal{H}}S := \bigcup_{p \in S} T_p^{\mathcal{H}}S$$

is a distribution of dimension $k-1$ if and only if S is noncharacteristic. The following proposition shows that $T^{\mathcal{H}}S$ depends only on Δ and S .

Proposition 3.5. *The horizontal tangent bundle $T^{\mathcal{H}}S$ is intrinsic:*

$$T^{\mathcal{H}}S = \Delta \bigcap TS$$

in the sense that $T_p^{\mathcal{H}}S = \Delta_p \bigcap T_p S$ for any $p \in S$.

Proof. Let g be any orthogonal extension of g_c . For $p \in S$, first let $v \in T_p^{\mathcal{H}}S$. Thus $v \in \Delta_p$ and from Definition 3.3 we have

$$\begin{aligned} g(v, n^g(p)) &= g_c(v, n^{\mathcal{H}}(p)) + g(v, n'(p)) \\ &= 0 + 0 = 0, \end{aligned}$$

where $n'(p) = n^g(p) - n^{\mathcal{H}}(p)$ is orthogonal to v since g is an orthogonal extension. The last formulas implies that $v \in T_p S$. So $T_p^{\mathcal{H}}S \subset \Delta_p \bigcap T_p S$. $\Delta_p \bigcap T_p S \subset T_p^{\mathcal{H}}S$ follows from the same argument. $\square$

Remark 3.6. Therefore $T^{\mathcal{H}}S$ is the projection of TS onto Δ . In the case of Carnot groups of step two, if $p \in S$ is not a characteristic point, $T_p^{\mathcal{H}}S$ has obvious geometric meaning: $T_p^{\mathcal{H}}S$ is the projection on the horizontal bundle of the Lie algebra of the tangent group which is the blowup set of S with respect to the natural dilations³, see [12]. This is the reason why we call $T_p^{\mathcal{H}}S$ horizontal tangent plane.

If $p \in S \setminus \Sigma_S$, let $\mathcal{V}(p) = \frac{n^{\mathcal{H}}(p)}{|n^{\mathcal{H}}(p)|}$ be the unit horizontal normal. Then by Proposition 3, $\mathcal{V}$ is intrinsic: independent of the choice of orthogonal extension, since $\Delta_p = T_p^{\mathcal{H}}S \oplus^{g_c} \mathcal{V}(p)$.

³That is the limit set of $S_{p,s} := \delta_{\frac{1}{s}}(p^{-1}S)$ under suitable topology as $s \rightarrow 0$.

A natural question is that whether there exist sufficiently many horizontal curves in hypersurfaces such that the intrinsic Carnot-Carathéodory metric can be defined. We will not pursue the general case. But we will prove that for contact manifolds of dimension more than three, there exists at least one smooth horizontal curve connecting two given points in a connected hypersurface S if Σ_S does not contain any immersed submanifold, and so the intrinsic Carnot-Carathéodory metric can be defined. This is the case if S is noncharacteristic or with isolated characteristic points. Note that even in the contact case, the horizontal connectivity in hypersurfaces is overlooked. Some authors asserted that there are few horizontal curves in hypersurfaces, even though these hypersurfaces are noncharacteristic, see e.g. [10] (p.485) where the authors wrote: “... Notice however that a H-regular hypersurface contains very few H-rectifiable curves; in particular we cannot define a geodesic distance on a H-rectifiable hypersurface...”⁴.

To prove Theorem 1.1, we first introduce the notion of the orbit of a family of vector fields.

Definition 3.7 (orbits of a family of vector fields). Let M be a connected smooth manifold and let $\mathcal{F}$ be any family of smooth vector fields globally defined on M . We define the orbit of a point $p \in M$ of this family as the set of points of M reachable piecewise by trajectories of vector fields in the family, that is,

$$\mathcal{O}_p := \{\exp(t_n f_k) \circ \cdots \circ \exp(t_1 f_1) \circ p \mid t_i \in \mathbb{R}, f_i \in \mathcal{F}, n \in \mathbb{N}\}$$

where $\exp(tf)(p)$ denotes the flow of the vector field f through p , i.e. the curve $\gamma(t)$ in M such that

$$\begin{cases} \dot{\gamma}(t) = f(\gamma(t)) \\ \gamma(0) = p \end{cases}.$$

Of course, if some of our vector fields are not complete then we consider only such $t_1, \dots, t_n$ for which the above expression has sense. It is clear that the relation: “ q belongs to the orbit of p ” is an equivalence relation on M and thus M is the disjoint union of orbits (equivalence classes).

The following orbit theorem is due to Sussmann (also Nagano), see [26].

Theorem 3.8 (Orbit Theorem, Nagano-Sussmann). *Let $\mathcal{F}$ be as above and let $p \in M$. Then:*

- (1) $\mathcal{O}_p$ is a connected immersion submanifold of M .
- (2) $T_q \mathcal{O}_p = \text{span}\{(P_* f)(q) \mid P \in \mathcal{P}, f \in \mathcal{F}\}$, $q \in \mathcal{O}_p$ where $\mathcal{P}$ is the group of diffeomorphisms of M generated by flows in $\mathcal{F}$:

$$\mathcal{P} = \{\exp(t_n f_k) \circ \cdots \circ \exp(t_1 f_1) \mid t_i \in \mathbb{R}, f_i \in \mathcal{F}, n \in \mathbb{N}\} \subset \text{Diff}(M)$$

and P_* is the differential map of P .

Remark 3.9. From Theorem 3.8, two simple but very useful observations are in order.

- (1) First of all, if $f \in \mathcal{F}$, then $f(q) \in T_q \mathcal{O}_p$ for all $q \in \mathcal{O}_p$. Indeed, the trajectory $\exp(tf)(q)$ belongs to the orbit $\mathcal{O}_p$, thus its velocity vector $f(q)$ is in the tangent space $T_q \mathcal{O}_p$.
- (2) Further, if $f_1, f_2 \in \mathcal{F}$, then $[f_1, f_2](q) \in T_q \mathcal{O}_p$ for all $q \in \mathcal{O}_p$. This follows since the vector $[f_1, f_2](q)$ is tangent to the trajectory $\exp(-tf_2) \circ \exp(-tf_1) \circ \exp(tf_2) \circ \exp(tf_1)(q) \in \mathcal{O}_p$. We go on and consider Lie brackets of arbitrarily high order

$$[f_1, [\cdots [f_{n-1}, f_n] \cdots]](q)$$

as tangent vectors to $\mathcal{O}_p$ if $f_i \in \mathcal{F}$ and $q \in \mathcal{O}_p$.

⁴Smooth noncharacteristic hypersurfaces are, from the definition, H-regular hypersurfaces; the inverse is not true in general. H-rectifiable curves are just absolutely continuous horizontal curves.

These considerations can be summarized in terms of Lie algebra of vector fields generated by $\mathcal{F}$:

$$\text{Lie}\mathcal{F} := \text{span}\{[f_1, [\cdots [f_{n-1}, f_n] \cdots]] \mid f_i \in \mathcal{F}, n \in \mathbb{N}\} \subset (\Gamma(TM)),$$

and its evaluation at a point $q \in M$:

$$\text{Lie}_q\mathcal{F} = \{V(q) \mid V \in \text{Lie}\mathcal{F}\} \subset T_qM.$$

We obtain the following statement.

Corollary 3.10.

$$\text{Lie}_q\mathcal{F} \subset T_q\mathcal{O}_p$$

for all $q \in \mathcal{O}_p$.

We note that the Chow connectivity theorem follows immediately from Corollary 3.10.

Now we return to consider the horizontal connectivity in hypersurfaces in sub-Riemannian manifolds. We will need the notion of a horizontal immersed submanifold with respect to a distribution.

Definition 3.11. Let M be a smooth manifold and let Δ be a distribution on M . An immersed submanifold $i : N \rightarrow M$ of M is horizontal with respect to Δ if $i_*(T_pN) \subset \Delta_p$ for any $p \in N$.

For codimension one distribution we have

Theorem 3.12. *Let M be a smooth connected manifold of dimension m equipped with a smooth distribution Δ of dimension $m - 1$. If M does not admit horizontal immersed submanifolds of dimension $m - 1$, then given two points p, q in M , there exists at least one piecewise smooth horizontal (with respect to Δ) curve connecting them (in fact Δ is equiregular).*

Proof. Let $\mathcal{F}$ be the set of all smooth vector fields tangent to Δ , that is,

$$\mathcal{F} = \{f \in \Gamma(TM) \mid f(p) \in \Delta_p \text{ for any } p \in M\}.$$

For any $p \in M$ we claim that the orbit $\mathcal{O}_p$ of p of the family $\mathcal{F}$ is of full dimension and thus $\mathcal{O}_p$ is an open set of M .

In fact, if not, then $\dim(\mathcal{O}_p) \leq m - 1$ where $\dim(\mathcal{O}_p)$ denotes the dimension of $\mathcal{O}_p$. On the other hand by Corollary 3.10, we have $\Delta_q \subset T_q\mathcal{O}_p$ for any $q \in \mathcal{O}_p$ and so the dimension of $\mathcal{O}_p$ is not less than $\dim(\Delta) = m - 1$. Thus, $\dim(\mathcal{O}_p) = m - 1$ and $T_q\mathcal{O}_p = \Delta_q$ for any $q \in \mathcal{O}_p$. So $\mathcal{O}_p$ is a horizontal immersed submanifold of dimension $m - 1$. This contradicts with the assumption.

Since M is connected and M is the union of all orbits, we have $M = \mathcal{O}_p$ for any $p \in M$. $\square$

Lemma 3.13. *Let M be a smooth orientable contact manifold of dimension $2n+1$ with a contact form η . Then M does not admit horizontal (with respect to $\Delta = \ker(\eta)$) immersion submanifolds of dimension more than n .*

Proof. This is a well known fact. For the readers' convenience and completeness we give a proof.

Let $i : N \rightarrow M$ be a horizontal immersed submanifold of M and let $p \in N$. By definition, $\dim(i_*(T_pN)) = \dim(T_pN)$ and for any $v_1, v_2 \in i_*(T_pN)$,

$$\omega_{i(p)}(v_1, v_2) = 0 \tag{3.1}$$

where $\omega = d\eta$. Since $\omega_{i(p)}$ is a symplectic form on the horizontal space Δ_p , from the non-degeneracy of $\omega_{i(p)}$ we have that

$$\dim((i_*(T_pN))^\perp) + \dim(i_*(T_pN)) = 2n$$

where $(i_*(T_p N))^\perp$ is the symplectic orthogonal subspace of $i_*(T_p N)$. (3.1) is equivalent to $i_*(T_p N) \subset (i_*(T_p N))^\perp$. Thus

$$2 \dim i_*(T_p N) \leq \dim((i_*(T_p N))^\perp) + \dim(i_*(T_p N)) = 2n.$$

□

Now we are in the position to prove Theorem 1.1.

Proof of Theorem 1.1. Since $2n - 1 > n$, the statement follows directly from Theorem 3.12 and Lemma 3.13 if S is noncharacteristic. If S possesses characteristic points, let $\mathcal{F}_S^\Delta$ be the set of all smooth vector fields tangent to $T^\mathcal{H}S$, that is,

$$\mathcal{F}_S^\Delta := \{f \in \Gamma(TS) \mid f(p) \in T_p^\mathcal{H}S \text{ for any } p \in S\}.$$

Let $\mathcal{O}_p$ be an orbit of $p \in S$ of the family $\mathcal{F}_S^\Delta$. By Theorem 3.8 $\mathcal{O}_p$ is an immersed submanifold. From the assumption that Σ_S does not contain immersed submanifolds we conclude that there exists at least one point $q \in \mathcal{O}_p$ which is not a characteristic point. By Corollary 3.10 we have that $\mathcal{F}_S^\Delta(q) \subset T_q(\mathcal{O}_p)$ and the dimension of $\mathcal{O}_p$ is not less than $2n - 1 = \dim(\mathcal{F}_S^\Delta(q))$. Thus if $\dim(\mathcal{O}_p) = 2n - 1$, then $\mathcal{O}_p$ is a horizontal immersion submanifold of S (and M). This contradicts with Lemma 3.13 since $n > 1$. So $\mathcal{O}_p$ is of full dimension and it is an open set of S . The assertion follows from the connectedness of S . □

Example 3.14 (the gauge ball in $\mathbb{H}^n$ with $n > 1$). Let $\mathbb{S}^n = \{p \in \mathbb{R}^{2n+1} \mid \|p\| = 1\}$ where $\|p\| := ((|x|^2 + |y|^2)^2 + |t|^2)^{\frac{1}{4}}$ is the gauge norm in $\mathbb{H}^n$. $\mathbb{S}^n$ is called the gauge ball centered in the origin. It is trivial to check that the metric induced by the gauge norm is left-invariant and 1-homogeneous with respect to natural dilations in $\mathbb{H}^n$. By direct computation the characteristic set $\Sigma_{\mathbb{S}^n}$ of $\mathbb{S}^n$ consists of only two points:

$$\Sigma_{\mathbb{S}^n} = \{(0, 0, 1), (0, 0, -1)\}.$$

By Theorem 1.1, we see that the induced Carnot-Carathéodory metric $\mathbb{S}^n$ can be defined.

Example 3.15 (hyperplanes in $\mathbb{H}^n$ with $n > 1$). Let $n > 1$. The vertical hyperplane $L_i = \{(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_n, y, t) \in \mathbb{R}^{2n+1}\}$ of $\mathbb{H}^n$ is a Lie subgroup of $\mathbb{H}^n$ with the induced group law (that is the restriction to L_i of the group law of $\mathbb{H}^n$) and Lie algebra $\mathcal{L}_i := \overline{V}_1 \oplus \overline{V}_2$ where $\overline{V}_1 = \text{span}\{X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n, X_{n+1}, \dots, X_{n+i-1}, \overline{X}_{n+i}, X_{n+i+1}, \dots, X_{2n}\}$ and $\overline{V}_2 = \text{span}\{T\}$ where $\overline{X}_{n+i} := \frac{\partial}{\partial y_i}$ and X_i, T as in Example 2.3). L_i is noncharacteristic and is a Carnot group. It is easy to prove that the Carnot-Carathéodory metric d_{in} induced by $(\overline{V}_1, g_c^i)$ where g_c^i is the restriction of g_c to $\overline{V}_1$ and the restricted Carnot-Carathéodory metric d_{re} satisfies that

$$d_{re} \leq d_{in} \leq C d_{re}$$

where C is an absolute constant, see [17], and there are points $p, q \in L_i$ such that $d_{re}(p, q) < d_{in}(p, q)$.

The following example shows the condition of dimension more than three is unavoidable.

Example 3.16. In the simplest Heisenberg group $\mathbb{H}^1$, we consider horizontal curves in the horizontal hyperplane $L_t = \{(x, y, 0) \in \mathbb{R}^3\}$. Note that the point $(0, 0, 0)$ is the unique characteristic point in L_t . Let $\gamma(t) = (x(s), y(s), 0)$ be a horizontal curve in L_t . Then from (2.1) we have

$$x(s)\dot{y}(s) = y(s)\dot{x}(s)$$

and hence

$$y(s) = Cx(s)$$

for some positive constant C . Thus there are no horizontal curves in any hypersurface $S \subset L_1$ which does not contain the point $(0, 0, 0)$. The same argument shows that there are no horizontal curves in $L_x = \{(0, y, t) \in \mathbb{R}^3\}$ and in $L_y = \{x, 0, t\} \in \mathbb{R}^3\}$.

Remark 3.17. As in Euclidean geometry, $\mathbb{R}^3$ can serve as a model to the study of higher dimension, in the study of sub-Riemannian geometry $\mathbb{H}^1$ can also be seen as a model. But in the developing geometric measure theory in the setting of sub-Riemannian geometry, in particular for the notion of rectifiability (and possibly for co-area formulae) in Carnot groups, $\mathbb{H}^1$ may be an exception. We recall that in [10]–[13] B. Franchi, R. Serapioni and F. Serra Cassano have proposed a notion of rectifiability by introducing the notion of intrinsic regular hypersurfaces, and another notion which is a counterpart of Federer’s definition of rectifiability where the “model spaces” are replaced by Carnot groups is announced by S. Pauls in [24]. It seems that the Pauls’ notion is meaningless for $\mathbb{H}^1$. One reason is that the codimension one Lie subgroups L_x, L_y of $\mathbb{H}^1$ have no stratified structure. Another reason is the horizontal non-connectivity of hypersurfaces in $\mathbb{H}^1$ as shown in the last example. A very intriguing question arises: whether are the two notions of rectifiability for $\mathbb{H}^n (n > 1)$ equivalent in any reasonable sense?

4. HORIZONTAL CONNECTION, THE HORIZONTAL MEAN CURVATURE AND THE HORIZONTAL DIVERGENCE THEOREM

Definition 4.1 (horizontal connection). Let g be any orthogonal extension of g_c and let ∇ be the Levi-Civita connection with respect to g . We define the horizontal connection $\tilde{D}$ on Δ as

$$\begin{aligned} \tilde{D} : \Gamma(\Delta) \times \Gamma(\Delta) &\rightarrow \Gamma(\Delta) \\ \tilde{D}_X Y &= \sum_{i=1}^k g(\nabla_X Y, X_i) X_i \text{ for any } X, Y \in \Gamma(\Delta) \end{aligned}$$

where $\{X_1, \dots, X_k\}$ is an orthonormal basis of Δ .

Remark 4.2. The definition of $\tilde{D}$ is independent of the choice of orthonormal basis of Δ . In fact, let $\bar{X}_i = \sum_{j=1}^k a_{ij} X_j, i = 1, \dots, k$ be another orthonormal basis. Then (a_{ij}) is an orthonormal matrix (everywhere) and hence

$$\begin{aligned} \sum_{i=1}^k g(\nabla_X Y, \bar{X}_i) \bar{X}_i &= \sum_{j=1}^k \sum_{l=1}^k \sum_{i=1}^k a_{ij} a_{il} g(\nabla_X Y, X_j) X_l = \sum_{j=1}^k \sum_{l=1}^k \delta_{jl} g(\nabla_X Y, X_j) X_l \\ &= \sum_{j=1}^k g(\nabla_X Y, X_j) X_j \end{aligned}$$

Thus $\tilde{D}$ is well defined.

We call $\tilde{D}$ a “connection” because of the following fact.

Lemma 4.3. $\tilde{D}$ satisfies the following properties

- (1) $\tilde{D}_X Y$ is $\mathbb{R}$ –linear in both arguments,
- (2) $\tilde{D}_X Y$ is $C^\infty(M)$ –linear in the argument of X ,
- (3) the Leibniz rule holds:

$$\tilde{D}_X(fY) = (Xf)Y + f\tilde{D}_X Y \text{ for any } f \in C^\infty(M), X, Y \in \Gamma(\Delta)$$

(4) $\tilde{D}$ is compatible with respect to g_c , that is,

$$Xg_c(Y, Z) = g_c(\tilde{D}_X Y, Z) + g_c(Y, \tilde{D}_X Z) \text{ for any } X, Y, Z \in \Gamma(\Delta), \quad (4.1)$$

(5) the following “symmetry” property holds:

$$\tilde{D}_X Y - \tilde{D}_Y X = [X, Y]^{\mathcal{H}} \text{ for any } X, Y \in \Gamma(\Delta) \quad (4.2)$$

where $[X, Y]^{\mathcal{H}}$ is, by definition, the projection of $[X, Y]$ to Δ , that is,

$$[X, Y]^{\mathcal{H}} = \sum_{i=1}^k g([X, Y], X_i) X_i.$$

Proof. The proof is trivial. All follow directly from the definition of $\tilde{D}$, the compatibility and the symmetry of the Levi-Civita connection ∇ together with the fact that g is an orthogonal extension of g_c . $\square$

Remark 4.4. For any vector field X , the horizontal part $X^{\mathcal{H}}$ of X : $X^{\mathcal{H}} = \sum_{i=1}^k g(X, X_i) X_i$, is independent of any orthogonal extension g of g_c . In fact, since g is an orthogonal extension of g_c , the projection, with respect to the decomposition of direct summation, of a vector field to the horizontal bundle is the same as the projection of this vector field to the horizontal bundle, with respect to the orthogonal decomposition.

From (2) and (3) of Lemma 4.3 it is straight to verify that $\tilde{D}_X Y(p)$ depends only on $X(p)$ and the evaluations of Y in a neighborhood of p .

Any operator from $\Gamma(\Delta) \otimes \Gamma(\Delta) \rightarrow \Gamma(\Delta)$ satisfying (1), (2), (3) and (4) of Lemma 4.3 is called a *nonholonomic connection on Δ* .

Proposition 4.5. *The operator $\tilde{D}$ is independent of the choice of orthogonal extensions of g_c .*

Proof. From Lemma 4.3 we know that $\tilde{D}_U V = \sum_{j=1}^k U(V^j) X_j + \sum_{i=1}^k \sum_{j=1}^k U^i V^j \tilde{D}_{X_i} X_j$ for any $U = \sum_{i=1}^k U^i X_i$, $V = \sum_{j=1}^k V^j X_j \in \Gamma(\Delta)$. Thus $\tilde{D}$ is determined by the connection coefficients $\Gamma_{ij}^l = g_c(\tilde{D}_{X_i} X_j, X_l) = g(\nabla_{X_i} X_j, X_l)$. By the Cozhul’s formula, for any $i, j, l = 1, \dots, k$ we have

$$\begin{aligned} g(\nabla_{X_i} X_j, X_l) &= \frac{1}{2} \{ X_i g(X_j, X_l) + X_j g(X_l, X_i) - X_l g(X_i, X_j) \\ &\quad - g(X_i, [X_l, X_j]) - g(X_j, [X_i, X_l]) + g(X_l, [X_j, X_i]) \} \\ &= -\frac{1}{2} \{ g(X_i, [X_l, X_j]) + g(X_j, [X_i, X_l]) - g(X_l, [X_j, X_i]) \} \\ &= -\frac{1}{2} \{ g_c(X_i, [X_l, X_j]^{\mathcal{H}}) + g_c(X_j, [X_i, X_l]^{\mathcal{H}}) - g_c(X_l, [X_j, X_i]^{\mathcal{H}}) \}, \end{aligned}$$

since $\{X_1, \dots, X_k\}$ is an orthonormal basis of Δ with respect to g_c . The assertion follows from Remark 4.4. $\square$

Now we turn to the proof of Theorem 1.2. The definition of D in Theorem 1.2 is inspired by the proof of Proposition 4.5 and the fundamental theorem in Riemannian geometry.

Proof of Theorem 1.2. In this proof, to simplify the notations we use the Einstein summation convention: if in any term the same index name appears twice, as both an upper and a lower index, that term is assumed to be summed over all possible values of that index (from 1 to k). We first prove that D is independent of the choice of orthonormal basis of Δ . Let $\{\tilde{X}_1, \dots, \tilde{X}_k\}$ be another orthonormal basis of Δ , where $\tilde{X}_i = a_i^j X_j$, $i = 1, \dots, k$ and $A = (a_i^j)$ is an orthogonal matrix (everywhere). For any $U = \tilde{U}^j \tilde{X}_j = U^j X_j$, $V = \tilde{V}^i \tilde{X}_i = V^i X_i \in \Gamma(\Delta)$ where $\tilde{U}^j = b_r^j U^r$,

$\tilde{V}^i = b_s^i V^s$ and $B = (b_i^j)$ is the inverse matrix of A with $b_i^j = a_j^i$ for $i, j = 1, \dots, k$ since A is orthogonal, we try to compute

$$U(\tilde{V}^i)\tilde{X}_i + \tilde{U}^j\tilde{V}^i\tilde{\Gamma}_{ij}^l\tilde{X}_l = \mathbf{I} + \mathbf{II}$$

where $\tilde{\Gamma}_{ij}^l = -\frac{1}{2}\{g_c(\tilde{X}_i, [\tilde{X}_j, \tilde{X}_l]^{\mathcal{H}}) + g_c(\tilde{X}_j, [\tilde{X}_l, \tilde{X}_i]^{\mathcal{H}}) - g_c(\tilde{X}_l, [\tilde{X}_i, \tilde{X}_j]^{\mathcal{H}})\}$. Since

$$[\tilde{X}_j, \tilde{X}_l] = a_j^r a_l^h [X_r, X_h] + a_j^r X_r (a_l^h) X_h - a_l^h X_h (a_j^r) X_r,$$

we have

$$[\tilde{X}_j, \tilde{X}_l]^{\mathcal{H}} = a_j^r a_l^h [X_r, X_h]^{\mathcal{H}} + a_j^r X_r (a_l^h) X_h - a_l^h X_h (a_j^r) X_r$$

and hence

$$g_c(\tilde{X}_i, [\tilde{X}_j, \tilde{X}_l]^{\mathcal{H}}) = a_i^s a_j^r a_l^h g_c(X_s, [X_r, X_h]^{\mathcal{H}}) + \sum_{s=1}^k a_i^s a_j^r X_r (a_l^s) - \sum_{s=1}^k a_i^s a_l^h X_h (a_j^s),$$

using the same arguments to other terms, and from $b_i^j a_l^i = \delta_l^j$, $a_l^i a_i^j = \delta_i^j$ for $i, j = 1, \dots, k$, we deduce

$$\begin{aligned} \mathbf{II} = & -\frac{1}{2} \sum_{h=1}^k V^s U^r \{g_c(X_s, [X_r, X_h]^{\mathcal{H}}) + g_c(X_r, [X_h, X_s]^{\mathcal{H}}) - g_c(X_h, [X_s, X_r]^{\mathcal{H}})\} X_h \\ & -\frac{1}{2} \left\{ \sum_{s=1}^k \sum_{l=1}^k U^r V^s a_l^h X_r (a_l^s) X_h - \sum_{s=1}^k \sum_{h=1}^k U^r V^s b_r^j X_h (a_j^s) X_h - U^r V^s b_r^j X_s (a_j^h) X_h \right. \\ & \left. + U^r V^s b_s^i X_r (a_i^h) X_h - \sum_{r=1}^k \sum_{h=1}^k U^r V^s b_s^i X_h (a_i^r) X_h + \sum_{r=1}^k \sum_{l=1}^k U^r V^s a_l^h X_s (a_l^r) X_h \right\}. \end{aligned}$$

Renaming and rearranging some indices, then cancelling or adding some terms (using $X_l(a_r^j b_j^h) = 0$ and $a_l^r = b_r^l$ for any $l, r, h = 1, \dots, k$), we infer that

$$\mathbf{II} = U^r V^s \Gamma_{sr}^h X_h - U^r V^j a_i^l X_r (b_j^i) X_l.$$

Since

$$\begin{aligned} \mathbf{I} &= U(\tilde{V}^j)\tilde{X}_j = U^r X_r (b_i^j V^i) a_j^l X_l \\ &= U^r X_r (V^i) b_i^j a_j^l X_l + U^r V^i X_r (b_i^j) X_l \\ &= U(V^j) X_j + U^r V^i X_r (b_i^j) a_j^l X_l, \end{aligned}$$

we obtain

$$U(\tilde{V}^i)\tilde{X}_i + \tilde{U}^j\tilde{V}^i\tilde{\Gamma}_{ij}^l\tilde{X}_l = U(V^j) X_j + U^r V^s \Gamma_{sr}^h X_h.$$

Thus D is well defined.

From the definition of D we have for any $i, j, l = 1, \dots, k$,

$$\Gamma_{ij}^l = g_c(DX_j X_i, X_l), \quad (4.3)$$

$$\Gamma_{ji}^l + \Gamma_{li}^j = 0, \quad (4.4)$$

$$\Gamma_{ij}^l - \Gamma_{ji}^l = g_c([X_j, X_i]^{\mathcal{H}}, X_l). \quad (4.5)$$

Properties (1)-(3) are obvious. To prove (1.2), let $U = U^j X_j, V = V^i X_i, W = W^l X_l$. Then from (2) and (3) we have

$$\begin{aligned} g_c(D_V U, W) + g_c(U, D_V W) &= U^j W^l g_c(D_{X_j}(V^i X_i), X_l) + U^j V^i g_c(X_j, D_{X_i}(W^l X_l)) \\ &= U^j V^i W^l (g_c(D_{X_j} X_i, X_l) + g_c(X_j, D_{X_i} X_l)) + \sum_{i=1}^k W^i V(U^i) \\ &\quad + \sum_{j=1}^k U^j V(W^j) \\ &= U^j V^i W^l (g_c(D_{X_j} X_i, X_l) + g_c(X_j, D_{X_i} X_l)) + V g_c(W, U). \end{aligned}$$

Since by (4.3) and (4.4)

$$\begin{aligned} g_c(D_{X_j} X_i, X_l) + g_c(X_j, D_{X_i} X_l) &= \Gamma_{ji}^l + \Gamma_{lj}^i \\ &= 0 \end{aligned}$$

for any $i, j, l = 1, \dots, k$, we conclude that

$$g_c(D_V U, W) + g_c(U, D_V W) = V g_c(W, U).$$

To see (1.3), for any $l = 1, \dots, k$, $U = U^j X_j, V = V^i X_i \in \Gamma(\Delta)$, by (4.5) we have

$$\begin{aligned} g_c(D_U V - D_V U, X_l) &= U(V^l) - V(U^l) + U^j V^i (\Gamma_{ij}^l - \Gamma_{ji}^l) \\ &= U(V^l) - V(U^l) + U^j V^i g_c([X_j, X_i]^{\mathcal{H}}, X_l) \\ &= g_c([U, V]^{\mathcal{H}}, X_l). \end{aligned}$$

The uniqueness follows from an argument similar to the Riemannian case: first use (2) and (3) to obtain (1.1) where $\Gamma_{ij}^l = g_c(D_{X_j} X_i, X_l)$, then using (1.2) and (1.3) to get a formula of Cozhul type, $g_c(D_{X_j} X_i, X_l) = -\frac{1}{2} \{g_c(X_i, [X_l, X_j]^{\mathcal{H}}) + g_c(X_j, [X_i, X_l]^{\mathcal{H}}) - g_c(X_l, [X_j, X_i]^{\mathcal{H}})\}$. $\square$

Remark 4.6. Thus D is an intrinsic notion of the sub-Riemannian structure (Δ, g_c) . Here we must point out that the connection $\bar{D}$ obtained by projecting the Levi-Civita connection onto the horizontal bundle is first introduced by Schouten in [25]⁵, further developed by Vagner in [28]-[31] (see [16] for a survey), see also [32] and [33] where $\bar{D}$ is called “truncated connection”. But it seems that they did not realize the existence of D in Theorem 1.2 and its intrinsic nature (determination by (Δ, g_c)) of D , see p.202 in [33].

In Russian mathematicians’ papers (also E. Cartan’s, see [3], [2] and [18]) nonholonomic connection was mainly used to study “geodesics” (but not sub-Riemannian geodesics i.e. non-holonomic geodesics) and curvature of distributions in the setting of nonholonomic dynamic systems. Let $\gamma(s)$ be a smooth horizontal curve in M . We call γ a “geodesic” if $D_{\dot{\gamma}} \dot{\gamma} = 0$. The mechanical significance of such “geodesics” is that they characterize the trajectories of motion of a mechanical system with quadratic Lagrangian and linear constraints (say Δ), see [32] and [33] for details.

Example 4.7 (the horizontal connection in Carnot group). In a Carnot group G , since its Lie algebra $\mathcal{G}$ is graded, if we choose a system of left invariant vector fields $\{X_1, \dots, X_k\}$ as an orthogonal basis of the horizontal bundle $\Delta = V_1$, then the connection coefficients Γ_{ij}^l of the

⁵There the Riemannian metric g with respect to which the Levi-Civita connection was computed is not necessary to be an orthogonal extension of g_c . Thus in this case $\bar{D}$ is not intrinsic: it depends on g .

horizontal connection D are vanishing everywhere for any $i, j, l = 1, \dots, k$. For this basis, D has the simple form

$$D_U V = \sum_{i=1}^k U(V^i) X_i \text{ for any } U, V = \sum_{i=1}^k V^i X_i \in \Gamma(\Delta),$$

see [27] for details.

For completeness we introduce the notion of horizontal divergence.

Definition 4.8 (horizontal divergence). Let $X \in \Gamma(\Delta)$. The horizontal divergence $\operatorname{div}_{\mathcal{H}} X$ of X is defined by

$$\operatorname{div}_{\mathcal{H}} X = \sum_{i=1}^k g_c(D_{X_i} X, X_i)$$

Note that as shown in Remark 4.2, $\operatorname{div}_{\mathcal{H}} X$ is independent of the choice of orthonormal basis of Δ because of (2) in Lemma 4.3.

The following proposition, which in the case of Carnot groups are well known, follows immediately from the definitions.

Proposition 4.9. $\operatorname{div}_{\mathcal{H}} X = \operatorname{div} X$ for any $X \in \Gamma(\Delta)$ where $\operatorname{div} X$ is the usual divergence of X computed with respect to any orthogonal extension g of g_c .

It is interesting that D can be used to define the horizontal mean curvature of (noncharacteristic) hypersurfaces in (M, Δ, g_c) . In the rest of this paper, we discuss how this can be done. Roughly speaking, in sub-Riemannian (or nonholonomic) geometry horizontal connection, horizontal tangent connection, horizontal normal, horizontal second fundamental form and horizontal mean curvature are counterparts of Levi-Civita connection, tangent connection, Riemannian normal, second fundamental form and mean curvature in Riemannian geometry, respectively.

We assume, if without further notice, S is a smooth **noncharacteristic** hypersurface in a sub-Riemannian manifold (M, Δ, g_c) . Then $T^{\mathcal{H}} S$ is a subbundle of TS of dimension $k-1$. From the definition we have

Lemma 4.10. If $X, Y \in \Gamma(T^{\mathcal{H}} S)$, then $[X, Y]^{\mathcal{H}} \in \Gamma(T^{\mathcal{H}} S)$.

It is clear that any vector v in $T_p^{\mathcal{H}} S (p \in S)$ can be extended to a vector field in $T^{\mathcal{H}} S$ by first extending v to a vector field V in TS then projecting V to $T^{\mathcal{H}} S$, and any vector field V in $T^{\mathcal{H}} S$ can smoothly extended to a horizontal vector field in Δ by first extending V to a vector field $\bar{V}$ in TM then projecting $\bar{V}$ to Δ . Sometimes we will denote by the same symbol both the extended vector field and the original vector field, in particular when we have confirmed that the objects under consideration are independent of such extensions.

If X, Y are vector fields in $T^{\mathcal{H}} S$, we can extend them to horizontal vector fields $\bar{X}, \bar{Y}$ in M , apply the ambient derivative operator D , and then decompose at points of S to get

$$D_{\bar{X}} \bar{Y}(x) = (D_{\bar{X}} \bar{Y})^{\top}(x) + (D_{\bar{X}} \bar{Y})^{\perp}(x) \quad x \in S \quad (4.6)$$

where $(D_{\bar{X}} \bar{Y})^{\top}(x)$, $(D_{\bar{X}} \bar{Y})^{\perp}(x)$ are the projections of $D_{\bar{X}} \bar{Y}(x)$ onto $T_x^{\mathcal{H}} S$ and the direction of $\mathcal{V}(x)$ respectively, where $\mathcal{V}(x)$ is the unit horizontal normal, see Remark 3.6.

Definition 4.11 (the horizontal second fundamental form). Let X, Y be vector fields in $T^{\mathcal{H}}S$. We define

$$\text{II}(X, Y) := (D_{\overline{X}}\overline{Y})^{\perp}$$

where $\overline{X}, \overline{Y}$ are the arbitrarily extended horizontal vector fields of X, Y respectively, as the horizontal second fundamental form of S .

Theorem 4.12. *The horizontal second fundamental form $\text{II}(X, Y)$ is*

- (1) *independent of the extension of X and Y ;*
- (2) *bilinear over $C^{\infty}(S)$; and*
- (3) *symmetric in X and Y .*

The proof of Theorem 4.12 is very similar to the Riemannian case, see [27]. Note that (3) follows from Lemma 4.10 and the symmetry property (4.2) of D , as in the Riemannian geometry the symmetry of the second fundamental follows directly from the symmetry (torsion free) of the Levi-Civita connection. Theorem 4.12 in particular implies that $\text{II}(X, Y)(p)$ depends only on $X(p)$ and $Y(p)$.

Definition 4.13 (the horizontal tangent connection). We define the horizontal tangent connection

$$D^{\top} : \Gamma(T^{\mathcal{H}}S) \otimes \Gamma(T^{\mathcal{H}}S) \rightarrow \Gamma(T^{\mathcal{H}}S)$$

by

$$D_X^{\top}Y = (D_{\overline{X}}\overline{Y})^{\top},$$

where $\overline{X}, \overline{Y}$ are the arbitrarily extended horizontal vector fields of X, Y respectively.

Theorem 4.14. *The horizontal tangent connection $D^{\top}$ is well defined, that is, $D_X^{\top}Y$ is independent of the extension of X and Y . Moreover $D^{\top}$ satisfies (1)-(5) of Lemma 4.3 where Δ is replaced by $T^{\mathcal{H}}S$ and g_c is replaced by $g_c^{\mathcal{H}}$ which is the restriction of g_c to $T^{\mathcal{H}}S$.*

The proof of Theorem 4.14 is direct and trivial. From the preceding discussions, we know that in S there exists an intrinsic horizontal connection determined by the sub-Riemannian structure $(T^{\mathcal{H}}S, g_c^{\mathcal{H}})$, since

$$TS = T^{\mathcal{H}}S \oplus^{\overline{g}} (TS/T^{\mathcal{H}}S)$$

where $\overline{g}$ is the restriction of g to TS . From the definitions it is easy to see that $\overline{g}$ is the orthogonal extension of $g_c^{\mathcal{H}}$, even if $T^{\mathcal{H}}S$ may not satisfy the Chow condition. Thus the “external” connection $D^{\top}$ is equal to the intrinsic one by Theorem 4.14 and Theorem 1.2.

Definition 4.15 (horizontal tangent divergence). Let $X \in \Gamma(T^{\mathcal{H}}S)$. The horizontal tangent divergence $\text{div}_{\mathcal{H}}^{\top}X$ of X is defined by

$$\text{div}_{\mathcal{H}}^{\top}X = \sum_{i=1}^{k-1} g_c^{\mathcal{H}}(D_{\tau_i}^{\top}X, \tau_i)$$

where $\{\tau_1, \dots, \tau_{k-1}\}$ is a orthonormal basis of $T^{\mathcal{H}}S$ with respect to $g_c^{\mathcal{H}}$. For a horizontal vector field Y defined in S , that is, $Y \in \Gamma(\Delta|_S)$ where $\Delta|_S$ is the restriction of Δ to S , we also can define its horizontal tangent divergence

$$\text{div}_{\mathcal{H}}^{\top}Y = \sum_{i=1}^{k-1} g_c(D_{\tau_i}Y, \tau_i).$$

Similar to Proposition 4.9, we have

Proposition 4.16. *Let $Y \in \Gamma(T^{\mathcal{H}}S)$. Then*

$$\operatorname{div}_{\mathcal{H}}^{\top} Y = \operatorname{div}_S Y$$

where $\operatorname{div}_S Y$ is the Riemannian tangent divergence of Y on S computed with respect to any orthogonal extension g of g_c .

Definition 4.17. The scalar horizontal second fundamental form h is the symmetric bilinear function on $T^{\mathcal{H}}S$ defined by

$$h(X, Y) = g_c(\Pi(X, Y), \mathcal{V}).$$

That is $\Pi(X, Y) = h(X, Y)\mathcal{V}$. Thus h uniquely determines an endomorphism of $T^{\mathcal{H}}S$, say A , that is,

$$g_c(AX, Y) = h(X, Y) \quad \text{for all } X, Y \in \Gamma(T^{\mathcal{H}}S).$$

A is self-adjoint and we call A the horizontal shape operator of S .

For any $p \in S$, A gives a symmetric linear map $A_p : T_p^{\mathcal{H}}S \rightarrow T_p^{\mathcal{H}}S$. Then by the symmetry of A_p , A_p has $k-1$ real eigenvalues.

Definition 4.18 (the horizontal mean curvature). The $k-1$ eigenvalues of A_p , $\kappa_1, \dots, \kappa_{k-1}$, are called the horizontal principal curvatures at p and the corresponding eigenspaces are called horizontal principal directions. We define the horizontal mean curvature $H_X(p)$ at p the trace of A_p , that is, $H_X(p) = \sum_{i=1}^{k-1} \kappa_i$ and call the product of $\kappa_1, \dots, \kappa_{k-1}$ the horizontal Gaussian curvature at p .

It has been proven in [27] that in the case of Carnot groups our definition of the horizontal mean curvature coincides with that in [6] and [23].

Remark 4.19. Since D , $T^{\mathcal{H}}S$ and $\mathcal{V}$ are intrinsic, so is the notion of horizontal mean curvature. As in [6] and [23], the horizontal mean curvature can be defined only at noncharacteristic points⁶. However by Proposition 3.2, the horizontal mean curvature can be defined almost everywhere.

Example 4.20. As in Example 4.7, choose a system of left-invariant vector fields $\{X_1, \dots, X_k\}$ as an orthogonal basis of the horizontal bundle, then for any noncharacteristic hypersurface $S \subset G$, the horizontal mean curvature can be expressed as the form

$$H(p) = \sum_{i=1}^k X_i(\mathcal{V}^i)(p) \quad \text{for any } p \in S$$

where $\mathcal{V} = \sum_{i=1}^k \mathcal{V}^i X_i$ is the unit horizontal normal of S , see [27].

REFERENCES

- [1] A. Bellaïche, Tangent spaces in sub-Riemannian geometry, in *Sub-Riemannian Geometry*, 1-78, Prog. in Math., v.144, edited by A. Bellaïche and J. J. Risler, Birkhäuser, 1996.
- [2] Anthony M. Bloch, P. S. Krishnaprasad, Jerrold E. Marsden, Richard M. Murray, Nonholonomic mechanical systems with symmetry, Arch. Rational Mech. Anal. **136**(1996), 21-99.
- [3] E. Cartan. Sur la representation geometrique des syst'emes materiels non holonomes, Proc. Int. Congr. Math., vol. 4(1928), Bologna, 253-261.
- [4] W. L. Chow, über Systeme non linearen partiellen Differentialgleichungen erster Ordnung. Math. Ann. **117**(1939), 98-105.

⁶In [6] (see also [15]), if p is a characteristic point, the horizontal mean curvature at p is defined by $H(p) = \lim_{q \rightarrow p} H(q)$ if this limit exists. Since it is impossible to determine the size of the set of characteristic points at which the limits exist, this definition seems meaningless

- [5] D. Danielli, N. Garofalo, D. M. Nhien, Minimal surfaces, surfaces of constant mean curvature and isoperimetry in Carnot groups, 2001, preprint.
- [6] D. Danielli, N. Garofalo, D. M. Nhien, Non-doubling Ahlfors measures, perimeter measures, and the characterization of the trace spaces of Sobolev functions in Carnot-Carathéodory spaces. Memo. AMS. to appear.
- [7] M. Derridj, Un problème aux limites pour une classe d'opérateurs du second ordre hypoelliptiques, Ann. Inst. Fourier, Grenoble, **21**(1971), 99-148.
- [8] M. Derridj, Sur un théorème de traces, Ann. Inst. Fourier, Grenoble, **22**(1972), 73-83.
- [9] B. Franchi, R. Serapioni, F. Serra Cassano, Meyers-Serrin type theorems and relaxation of variational integrals depending on vector fields. Houston Math. J. **22**(1996), 858-889.
- [10] B. Franchi, R. Serapioni, F. Serra Cassano, Rectifiability and perimeter in the Heisenberg group, Math. Ann., **321**(2001), 479-531.
- [11] B. Franchi, R. Serapioni, F. Serra Cassano, Regular hypersurfaces, intrinsic perimeter and implicit functions theorem in Carnot groups, to appear on Comm. in Analysis and Geometry(2001), available from <http://cvgmt.sns.it>.
- [12] B. Franchi, R. Serapioni, F. Serra Cassano, On the structure of finite perimeter sets in step 2 Carnot groups, 2001, preprint, available from <http://cvgmt.sns.it>.
- [13] B. Franchi, R. Serapioni, F. Serra Cassano, Rectifiability and perimeter in step 2 groups, 2001, preprint, available from <http://cvgmt.sns.it>.
- [14] N. Garofalo, D. M. Nhieu, Isoperimetric and Sobolev inequalities for Carnot-Carathéodory spaces and the existence of minimal surfaces, Comm. Pure Appl. Math. **49**(1996), 1081-1144.
- [15] N. Garofalo, S. D. Pauls, The Bernstein problem in the Heisenberg group, 2002, preprint, available from: <http://cvgmt.sns.it>.
- [16] E. M. Gorbatenko, Differential geometry of nonholonomic manifolds (after V.V.Vagner), Geom. Sbornik, 31-43, Tomsk University, Tomsk, 1985.
- [17] M. Gromov, Carnot-Carathéodory spaces seen from within, in *Sub-Riemannian Geometry*, 79-323, Prog. in Math., v.144, edited by A. Bellaïche and J. J. Risler, Birkhäuser, 1996.
- [18] J. Koiller, P. R. Rodrigues, P. Pitanga, Non-holonomic connections following Élie Cartan, An. Acad. bras. Cienc. **7**(2001) 165-190.
- [19] U. Koschorke, *Vector fields and other vector bundle morphisms-a singularity approach*. Lecture notes in Mathematics 847, Springer, Berlin, 1981.
- [20] V. Magnani, *Elements of geometric measure theory on sub-Riemannian groups*, Ph.D. thesis, Scuola Normale Superiore, Pisa, 2002.
- [21] V. Magnani, A blow-up theorem for regular hypersurfaces on nilpotent groups, Manuscripta Math., **110**(2003), 55-76.
- [22] R. Montgomery, *A Tour of Subriemannian Geometry, Their Geodesics and Applications*, Mathematical Surveys and Monographs, vol. 91, 2002.
- [23] S. D. Pauls, Nonparametric minimal surfaces in the Heisenberg group, to appear in Geometriae Dedicata.
- [24] S.D. Pauls, A notion of rectifiability modelled on Carnot groups, preprint 2001.
- [25] J. A. Schouten, On nonholonomic connections, Proc. Amsterdam Nederl. Akad. Wetensch., Ser. A, **31**(1928), 291-299, Jbuch 54, 758.
- [26] H. J. Sussmann, Orbits of families of vector fields and integrability of distributions, Trans. Amer. Math. Soc. **80**(1973), 171-188.
- [27] K. H. Tan, X. P. Yang, Horizontal connection and horizontal mean curvature in Carnot groups, preprint, 2003.
- [28] V. V. Vagner, Differential geometry of nonholonomic manifolds, Tr. Semin. Vektorn. Tenzorn. Anal. **213**(1935), 269-314(Russian).
- [29] V. V. Vagner, Differential geometry of nonholonomic manifolds, In: *The VIII-th International Competition for the N. I. Lobatschewski Prize Report*. The Kazan Physico-Mathematical Society, Kazan, 1940.
- [30] V. V. Vagner, A geometric interpretation of nonholonomic dynamical systems, Tr. Semin. Vektorn. Tenzorn. Anal. **5**(1941), 301-327.
- [31] V. V. Vagner, Geometria del calcolo delle variazioni, vol. 2 C.I.M.E., Roma.
- [32] A. M. Vershik, L. D. Faddeev, Lagrange mechanics in an invariant form, In: Probl. Theor. Phys. Vol.II, 129-141. English transl.: Sel. Math. Sov., **1**(1981), 339-350.
- [33] A. M. Vershik, V. Ya. Gershkovich, Nonholomic problems and the theory of distributions, Acta. Appl. Math., **12**, 181-209, 1988.

DEPARTMENT OF APPLIED MATHEMATICS, NANJING UNIVERSITY OF SCIENCE AND TECHNOLOGY, 210094,
NANJING, JIANGSU, P. R. CHINA

E-mail address: tankanghai2000@yahoo.com.cn

E-mail address: yangxp@mail.njust.edu.cn